

XJY-707

PRODUCT

Product Name:

Hydride Terminated Polydimethylsiloxanes

Structural Formula:

$$H_3C$$
 H_3C
 O
 CH_3
 O
 M
 CH_3
 O
 M
 CH_3
 O
 M
 CH_3


TYPICAL PROPERTIES Specification 707-10 707-015 707-030 707-050 707-075 707-30 Index Appearance colorless clear liquid Content of 0.27-0.30% 0.030-0.034% 0.048-0.052% 0.073-0.078% 0.09-0.11% 0.015-0.017% Hydrogen (%) Viscosity 24-40 100-150 50-80 30-50 4-7 250-350 (25°C, mm²/s) Volatile matter ≤10% ≤1.5% ≤10% ≤5% ≤3% or ≤1% ≤6%

Viscosity can be customized as customer's request.

APPLICATIONS

XJY-707 is soluble in various organic solvents such as aromatic hydrocarbons/petroleum hydrocarbons. It is insoluble in water and is a widely used silicone intermediate. It is usually used as a silicone end-capping agent. Due to containing reactive Si-H groups in the molecular structure, XJY-707 can be used in the synthesis of copolymer macromolecule by hydrosilylation. The synthesis of copolymer macromolecule can be made into a series of reactive silicone oil.

- As a chain extender for adding liquid silicone rubber, adding a crosslinking agent for forming a hot vulcanized silicone rubber;
- Used as a waterproofing agent and softener for natural and synthetic fabrics;
- Special silicone surfactant;
- Used in manufacture of a variety of end group reactive linear modified silicone oil;
- The key intermediates of block copolymerization reaction;
- Used for modification of the reaction, such as plastic resin;



PACKAGING, STORAGE AND TRANSPORTATION

XJY-707 is available in 20kg pails and 190kg plastic drums. Packaging options are available upon request.

This product stored at or below 25°C in the original unopened containers. Transported as non-dangerous goods.